



# Progress, challenges, and the role of public engagement to improve tick-borne disease literacy

Joyce M Sakamoto<sup>1,2</sup>

Vector-borne diseases have increased worldwide, facilitated by globalization and variations in climate. Tick and tick-borne disease researchers, veterinarians, medical practitioners, and public health specialists are working to share their expertise on tick ecology, disease transmission, diagnostics, and treatment in order to control tick-borne epidemics and potential pandemics. This review will be a brief overview of the current status of tick-borne diseases, challenges on the scientific and public fronts, and the role of public engagement in improving citizen education within the context of ticks and tick-borne disease research.

## Addresses

<sup>1</sup> Department of Entomology, The Pennsylvania State University, University Park, PA 16802 United States

<sup>2</sup> Center for Infectious Disease Dynamics, The Pennsylvania State University, University Park, PA 16802 United States

Corresponding author: Sakamoto, Joyce M ([jms1198@psu.edu](mailto:jms1198@psu.edu))

**Current Opinion in Insect Science** 2018, 28:81–89

This review comes from a themed issue on **Vectors and medical and veterinary entomology**

Edited by **Jason Rasgon**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 26th May 2018

<https://doi.org/10.1016/j.cois.2018.05.011>

2214-5745/© 2018 Elsevier Inc. All rights reserved.

## Introduction

Humans live in a globally connected world, where activities in one part of the world may have impacts on other distantly connected part. Fluctuations in climate have been correlated with changes in vector biology and increased vector-borne diseases at local and global scales [1–3,4<sup>•</sup>]. Although tick-borne diseases (TBD) may not be as dramatically impacted by sudden short-term changes in temperature as mosquito-borne diseases, warmer winters have been correlated with invasion and establishment of certain tick vector species into northern latitudes [3,4<sup>•</sup>,5].

Scientists, medical and veterinary practitioners, and public health specialists strive to understand vector biology and disease dynamics, design better diagnostic tools, and develop better treatments for vector-borne diseases on behalf of the public. TBD experts therefore have a

responsibility to disseminate their findings and recommendations to the public in a clear, concise manner that can be readily understood. This review provides an overview of the current status of research on ticks and TBD, challenges facing TBD experts, and the importance of active, bidirectional engagement with the public.

## Ticks

Ticks are obligate bloodfeeding arachnids in the order Acari [6]. Fossil evidence from Baltic amber suggest ticks have been parasitizing vertebrates since the Cretaceous (ca 99 MYA) [7,8]. Although known for their role as vectors of pathogens, ticks themselves are considered noxious pests whose bites can cause irritations, anemia, localized or systemic allergic reactions (including meat allergies), as well as more serious issues such as tick paralysis [9–11].

At the time of this writing, there are 722 described species of hard ticks (Ixodidae) and 208 species of soft ticks (Argasidae) [6]. Approximately 5% of known Ixodidae and 2% of Argasidae are known vectors of zoonotic diseases [12]. Since at least 38% of all tick species are known to bite humans (though not necessarily the intended hosts), it is likely that new TBD will continue to be identified [12,13<sup>••</sup>]. As human populations grow, broader encroachment and fragmentation of wild habitats may increase contact with pathogen-laden ticks [14].

## Global tick-borne diseases

TBD incidence is increasing at an alarming rate, signaling the need for a more proactive stance to prevent outbreaks of pandemic proportions [15]. While *Borrelia burgdorferi* (the causal agent of Lyme borreliosis) is by far the most prevalent tick-borne pathogen in the temperate countries of the world, other emerging or newly discovered tick-borne pathogens (many of which are potentially fatal) are also being described annually [4<sup>••</sup>,16–18]. Worldwide there are two TBD of pandemic or epidemic scale recognized by the World Health Organization (Crimean-Congo Hemorrhagic Fever and Tularemia [<http://www.who.int/emergencies/diseases/en/>]). Rocky Mountain Spotted Fever, an emergent TBD that is spreading throughout the Americas, can be transmitted by multiple tick species [19<sup>•</sup>]. Factors contributing to increased TBD incidence include habitat fragmentation, global travel, importation of animals, changing environmental conditions, and shifts in host populations in response to light pollution, habitat removal or encroachment, or altered migration behaviors of reservoir hosts) [4<sup>••</sup>,14,20–24].

While some countries have dedicated significant scientific, political, and/or monetary resources toward TBD research, other countries lack the facilities or funds for monitoring and combatting TBD, or do not consider TBD as top priorities for public health, particularly when they are being impacted by other vector-borne diseases [25]. Nevertheless, the impacts of TBD in poorer countries can exacerbate already-existing socioeconomic inequalities [25,26]. In a globally connected world, public health concerns in one country are no longer isolated and the ‘every country for itself’ attitude of the past no longer holds true. Global exports, exotic pet trade, and human travel have all contributed to introduction of novel vector species and their pathogens into nonnative areas [23,24,27]. Excluding global human traffic and trade, many migrating wild species (such as birds) travel between tropical and arctic biorealm and have been implicated as vehicles for invading tick species as far north as the arctic circle [28\*,29–31].

### Ticks and tick-borne disease in USA

Although 40 of 84 described species of ticks in the contiguous United States are known to bite humans, eight species are responsible for the majority of TBD [13\*\*,32,33]. These few species represent multiple taxa, yet the common characteristics that make them such effective vectors is the generality of host preference and the behavior of actively seeking out hosts. In contrast, many species of ticks are nidicolous, preferring to remain within the nest or burrows of their hosts for the entirety of their lifecycles [34]. However, if their preferred host is no longer available, some nidicolous species will readily seek non-typical hosts [35,36]. This can result in sudden outbreaks of diseases when an unsuspecting human comes in contact with a hungry tick (e.g. tickborne relapsing fever transmitted by the soft tick *Ornithodoros hermsii* feeding on rodents and humans co-inhabiting cabins) [35].

Between 2004 and 2016, reported cases of TBD have more than doubled [37]. Lyme disease (LD) (principally caused by *Borrelia burgdorferi sensu stricto* in the United States) accounted for 402,502 of the 642,602 total reported vector-borne (62.6%), and 82% of all reported TBD cases [18]. It is estimated that 300,000 cases of LD are diagnosed each year, although only 10% of these cases are actually reported to the Centers for Disease Control (CDC) [18]. LD was first described in the early 1970s, manifesting first as a juvenile form of arthritis in Lyme Connecticut, but until the incidence rates rose significantly, it was not a priority for public health [38,39]. Eventually, LD did garner enough national attention to inspire a Senate bill that increased federal funds for improving tick-borne disease-related activities (research, treatment, prevention, and education). In December of 2016, President Obama signed the *21st Century Cures Act*, which included authorization of the Department of Health and Human Services to support TBD research

and create a multidisciplinary ‘Tick-borne Disease Working Group’. Although national effort toward TBD control can be slow, there is at least hope that support is now available for research.

### Challenges on the scientific front

Vector biologists can play a key role in providing insight into environmental management to prevent tick infestation or disease transmission. Improving communication between vector researchers and other TBD-related fields would greatly improve disease surveillance. Some diagnostic facilities have an expert who can identify vector species using morphological and/or molecular characteristics (e.g. New York State Veterinary Diagnostic Laboratory).

There is a need for fast, affordable, and reliable multiplex diagnostic tools, and many new options are on the horizon. Whole genome sequencing technology has made it possible to identify novel tick-borne pathogens as they are encountered (such as *Borrelia mayonii*, Heartland and Bourbon viruses) [37]. While these diagnostic tools are promising, many are proprietary, making it difficult for those doing field vector surveillance to find a standardized diagnostic approach without sending off materials to a commercial laboratory.

### Challenges on the public front

When TBD specialists do not effectively communicate with the public, the public may not be aware of the research progress being made. As medical, veterinary, and scientific professions develop and test methods for combatting new or emerging tick-borne pathogens, the public is ultimately the intended beneficiary. Yet many TBD experts will refrain from public engagement, particularly when they feel an issue is politicized and that the intended audience does not want to listen.

Tick bite prevention education is not easy and may require repetition, persistence, and patience. In response to changes in climate and the eventual spread of LD, the Canadian government launched an aggressive campaign of tick and reservoir host surveillance, assessment of public TBD literacy, and public education in 2014 [3,40–42,43\*]. Over time they have observed that, despite overall increase in general awareness of LD, less than half the population was utilizing tick prevention strategies [44]. Humans can be slow to change health behavior, but communications on TBD risk need to be persistent about the importance of preventative measures for eventual change [44].

### TBD-literacy

The most effective strategies for tick bite prevention (and TBD transmission) are avoidance of high-risk areas, preventative approaches such as showers and tick checks, and proper application or usage of acaricides and/or

repellents. It is important to ensure that these strategies are clear to maximize the likelihood that the public will follow these strategies correctly and consistently.

One hurdle that a successful TBD prevention and management program needs to overcome is the level of TBD awareness of the average patient. Patients may not know how to prevent tick bites or identify symptoms beyond the not-always-present erythema migrans, the classic LD expanding annular rash (bullseye) [38]. Patients may hold misconceptions about TBD, often acquired through online browsing, social media, or television sources. Local or regional efforts to improve public education on tick bite prevention are sometimes available in more populated or well-funded areas, but these approaches often fail to effectively reach disadvantaged people, areas in which technology-based dissemination is not an option, or very rural or wilderness areas.

### Publicly available sources of TBD information

Passive dissemination of TBD knowledge has its role in providing information regarding the outcomes of data analysis and recommendations for high-risk areas. The Centers for Disease Control maintains several informative pages on their website ranging from recommendations on communicating risk, reporting recent TBD-related findings, as well as suggested approaches to communicating TBD risk for medical professionals [10] (<https://www.cdc.gov/ticks/>; <https://www.cdc.gov/nceizd/>; <https://www.cdc.gov/niosh/topics/tick-borne/resources.html>). The CDC site represents a wealth of information, but much of it is geared toward disseminating information to scientists, practitioners, or public health experts, or on best practices when communicating to the public. Although some pages are intended for the public, they can sometimes be difficult to navigate. Nevertheless, the information presented is usually up-to-date and cites information based on current or recently completed studies published in peer-reviewed journals.

Another publicly available, but not readily discoverable resource is available from the US Army. The Disease Epidemiology Division US Army Zoonotic Disease Report for 2015 was released by the Army Public Health Center in 2017 ([http://phc.amedd.army.mil/Periodical%20Library/ZDR\\_CY2015\\_v2.pdf](http://phc.amedd.army.mil/Periodical%20Library/ZDR_CY2015_v2.pdf)). These data reflect the One Health approach in that the data collected here include all zoonotic incidents from active-duty military personnel, incidents from non-active duty beneficiaries, the location of acquisition of zoonotic diseases, veterinary disease summaries on DoD-associated animals, and wildlife and vector surveillance results.

Several universities maintain TBD educational resource (e.g. University of Rhode Island's <http://www.tickencounter.org>). These sites balance information, levity, and effective visual aids to engage the nonscientist.

Some universities have developed apps, collect citizen-based data state- or nationwide (including photos of ticks), and present tick risk predictions relevant to season, tick species, and present regional TBD diseases. Tick-encounters.org uses compelling (and sometimes entertaining) multimedia presentations that appeal to the public sector ranging from computer graphics, mobile applications, resources on tick pathogen testing, and several educational how-to videos on pesticide application, tick checks, and tick removal.

### Dispelling misconceptions through engagement

Community engagement has been shown to be highly effective in providing citizens the opportunity to ask questions, voice concerns, and dispel fears about mosquito control strategies implemented by the World Mosquito Program (formerly Eliminate Dengue [45]). Similarly, TBD active engagement at the local or regional scales may encourage citizens to learn how to protect themselves, but also to dispel certain misconceptions about risk, symptom recognition, and prevention. For instance, TBD may be perceived as a disease of more rural environments, but in fact, people in urban areas may also be at risk if at least one vector and one zoonotic reservoir host are present [11,46]. Even though urban areas are mostly managed, many species of wildlife can adjust to and thrive in manmade habitats (parks, trails, gardens, attics, backyards, etc.) [11].

While it is assumed that people who engage in 'outdoor activities' are at risk, what constitutes 'outdoor activity' may not necessarily be as obvious as hunting, biking, or camping. Gardeners, pet owners, farmers, scouts, hunters, foresters, as well as citizens utilizing common green-spaces (e.g. parks) may be potentially exposed to tick habitats [44,47,48–50]. Those who provide support such as environmental health specialists or wilderness providers also need to be trained in being tick-vigilant [48,51].

### "One Health Initiative"

Globalization has connected the world in remarkable ways, but has also accelerated the spread of vector-borne diseases [37]. Each new or re-emerging infectious disease highlights the need for rapid disease prevention and control. This requires multidisciplinary communication between veterinary and medical health care workers, wildlife biologists, ecologists, anthropologists, and others. 'One Health' reflects the realization that emerging zoonotic diseases occur at the interfaces of animal–human–ecosystem [13].

Although veterinary and medical professionals have largely pioneered much of the effort, 'One Health Initiative' was intended to foster communication across multiple disciplines [52]. The value of One Health-based multidisciplinary communication between physicians and

veterinarians is evident in the example of a fatal disease outbreak is the Rocky Mountain Spotted Fever (RMSF) case in Mississippi [13<sup>••</sup>,53]. Two dogs misdiagnosed with ehrlichiosis died of RMSF, followed thereafter by their owner. Subsequently, there was an investigation and successful treatment of the remaining two dogs for RMSF. The surviving dogs had been infested with *Rhipicephalus sanguineus* (the brown dog tick), previously not considered a vector of RMSF [53]. This case identified *R. sanguineus* as a competent vector of RMSF and indicated that dogs can serve as sentinels for RMSF. Furthermore, if a dog is suspected of being infected with RMSF, both veterinary and medical professionals should report this to the appropriate medical authority (e.g. state public health agencies or the CDC) [53].

Human health is integrally tied to the health of domestic and wild animals. The importance of cross-communication between veterinary and medical professionals is evident, particularly in respect to zoonoses[13<sup>••</sup>,52,54]. Pet owners may have an increased risk of tick bites [47<sup>••</sup>]. Dogs can acquire ectoparasites when multiple dogs are in contact, or if the dogs are boarded in a kennel, and then bring the ticks home with them [55]. Similarly, cats that hunt outdoors may encounter wildlife (e.g. groundhogs, squirrels, chipmunks) and become exposed to zoonotic diseases [56–59]. More generally, humans and domesticated animals are at higher risk of acquiring tick-borne pathogens from reservoir hosts when living in close proximity to or in contact with wildlife [47<sup>••</sup>,57,60,61].

### Passive tick surveillance

Passive tick surveillance usually involves submissions by non-acarologists to experts at regional, state, or national level institutions. This requires doctors, veterinarians, or other citizens to submit tick samples for identification and/or testing for pathogens. The advantage to this approach is that it can cover a wide geographic range, providing a finer-scale resolution of the tick/TBD load [41,43<sup>•</sup>,62,63]. The main caveat is that associated data may be incomplete, the samples can be damaged during removal (losing important morphological characteristics), and there is an intrinsic host-association bias for ticks found on humans or domestic animals [64]. It is also time-consuming and effort-intensive to catalog and curate the database generated from the samples and the metadata [65,66]. Nevertheless, these data represent a measure of the risk of bites because the ticks are usually extracted from or associated with host tissue. Passive surveillance data can be complementary to active surveillance data, and can also be used to tailor experimental designs for downstream host-association studies using trapped, hunted, or road-killed animals, or combined with CO<sub>2</sub> or host volatile-modified trapping from nests or burrows [40,67–69].

Because animals may represent sentinels for zoonotic diseases, indirect mining of electronic pet health records

can be useful alternative for identifying increased risk of TBD through records of tick infestations [70]. This approach did not require intensive surveillance or molecular diagnostics and could serve as a cost-effective complement to an already existing tick surveillance program to identify pockets of tick activity that might be missed by passive surveillance alone. Collectively, tick from companion animals could be used to create a real-time map of TBD risk (e.g. <http://www.capcvet.org>).


### Novel public engagement approaches

Multi-disciplinary approaches to TBD education through citizen engagement can cultivate exciting, creative, and sometimes unusual methods (Figure 1). The Dutch National Institute for Public Health and the Environment (RIVM) has used a multimedia strategy (website, media, video game, leaflets) to target adults and children for TBD education [71,72]. Alone, these approaches have not been shown to significantly improve tick bite prevention, but together they work as complementary strategies to reinforce good preventative practices. Additionally, RIVM launched a website and an affiliated app (<https://www.tekenradar.nl/>) that allows citizens to record data associated with when they were bitten (age, photo of the tick, the bite, whether they have had Lyme disease before, whether or not they develop a bulls-eye rash) [73]. While citizens are responsible for initial symptom diagnosis and reporting, ticks can also be submitted for identification and testing by trained professionals [73]. Public submissions of tick photos for identification have also been used in Canada and in the USA [74].

Another nontraditional public engagement approach is the use of mass participation events (e.g. marathons) to engage a large group of people converging at a single location in contributing to a tick risk study [75<sup>••</sup>]. Mass participation events are great opportunities for large-scale citizen contributions in a localized area [75<sup>••</sup>]. In many ways, it has the advantages of a ‘bioblitz’, a group event in which scientists and citizens meet at a particular location for a brief period of time to capture and identify as many species in an area to estimate local biodiversity (<https://www.inaturalist.org/pages/bioblitz+guide>). However, the mass participation tick collection takes advantage of an already existing event to collect data and to educate the participants about tick checks and TBD.

Citizen science approach can be useful for fine-scale active tick surveillance [76]. Each citizen enrolled in the study monitored a small plot (e.g. their backyard) and provided repeated measures of tick load from multiple sources such as pets, gardens, or flags (using a white towel or cloth on vegetation) [76]. This fine-scale approach concurs with the finding by the CDC, which found that for residential properties, determining absolute numbers of infected ticks would provide a better estimate of risk, particularly if the degree of human use of

Figure 1



Sources of ticks from surveillance. Tick surveillance provides useful information that can be used to identify risk of tick bites and TBD traditional sources. Information can be from more traditional sources such as veterinary or medical hospitals or clinics, universities, or wildlife biologists. Other sources of ticks, host-association, or pathogen data may include tick testing companies, electronic veterinary record databases, mass participation events, or various public-based contributions (physical or photographic submissions, citizen scientists, etc).

the property is also taken into account [77]. This becomes a much more tenable undertaking once citizens are involved in the data collection process. In another study, Seifert *et al.* (2016) partnered with rural high school teachers to teach high school students about science, TBD, and tick surveillance [78<sup>\*</sup>]. This model could be extended toward other community-based groups such as elementary schools, school nurses, parent–teacher organizations, or clubs to train children and their parents about how to prevent tick bites.

### Suggestions for public engagement

One of the ways we can encourage TBD literacy is to cultivate more active community involvement in tick surveillance and provide control strategies that are easy to follow. There are many potential approaches that can be explored or implemented. Multiple social medial platforms, citizen science, groups, and formal science communication (scicomm) training of TBD experts collectively may expand the reach and efficacy of outreach education.

Approaches to improve TBD literacy need to go beyond the deficit model of public engagement (i.e. unidirectional dissemination of information). It is important to assess public familiarity with tick bite prevention, resistance to pesticide use, likelihood of compliance,

recognition of TBD symptoms, and necessity of consulting a physician or veterinarian. The public should also be directed to scientists who specialize in identifying ticks.

Social media can be highly effective at disseminating information rapidly, but requires careful planning and proper timing for maximum impact. The platform chosen for outreach needs to match the audience — different platforms appeal to different demographics [79,80]. Social media may also provide data on local or regional tick abundance and seasonality or report invasive tick species ([https://twitter.com/Contagion\\_Live/status/979056024119373826](https://twitter.com/Contagion_Live/status/979056024119373826)). However, strategic forethought and marketing expertise is advisable to avoid potential unwanted outcomes (e.g. ticks on poppy seed muffins, <https://twitter.com/CDCgov/status/993523011281145858>).

Citizen science can be used to facilitate public submission of data and allow interested citizens to be more actively involved in a scientific process that may benefit them. Citizen science-based projects allow scientists to increase the amount of data collected or handled that would otherwise be difficult to achieve. It is important, however, to ensure a bidirectional conversation between the public and researchers, providing regular updates on the progress of studies in which they are participating.



Formal training of TBD specialists in science communication is not normally part of graduate or postgraduate career experience, but perhaps it should be. TBD specialists need to be able to disseminate their data beyond the academic and professional spheres to journalists, policy makers, stakeholders, industry, but without the use of jargon [81<sup>•</sup>]. Developing positive bidirectional lines of communication between TBD experts and the media provides journalists with news on scientific progress and insight into what tick species is a vector of which disease, and gives scientists the opportunity to reach a large audience. TBD experts in turn need to listen and adjust management recommendations while taking into consideration the cultural, socio-economical, and physical limitations of citizens.

Best models for engagement may be dependent on the target audience. An effective engagement approach would be to firstly keep the message simple and jargon-free, secondly have easy-to-understand protocols, and lastly use visual aids. While there is definitely an appeal to incorporate next-gen technology (e.g. augmented, virtual, or mixed reality), the emphasis should be on the message and not on the 'glitz'.

## Conclusion

TBD researchers, practitioners, and experts have the advantage over many other members of the scientific community. The public loves to hate ticks and is therefore much more likely to engage in conversation about ticks, TBD, and ways to prevent tick bites. This makes public engagement that much easier, but taken a step further, public involvement can provide insight into activities and/or studies in which the average citizen can feel personally empowered as contributors to the scientific process.

Although there may be the perception amongst scientists that the public does not trust scientific research, this is a misconception and in fact the opposite is true [82]. Scientists are becoming acutely aware of the importance of effective science communication and this applies to TBD experts as well. It is the responsibility of those studying vector biology, vector-borne diseases, and public health to teach the public how to protect themselves, prevent or minimize vector-borne disease transmission, recognize symptoms that require immediate attention, and to record information that will facilitate correct diagnosis and treatment. Transitioning from simply providing information to actively engaging communities will ultimately improve our vector-borne disease education efforts.

The United States is a very large country, so while there are many great local or regional efforts, there is still a need for unifying TBD education efforts and integration of resources. Efforts at state or regional levels vary — in some states, dedicated staff members spend a great deal

of effort on vector control and on public outreach, while in others, the burden falls on academic institutions or public health departments. Perhaps with the passing of the 21st Century Cures Act and the CDC report on vector-borne disease on the rise, support for improved TBD public education is just on the horizon [37,83<sup>••</sup>].

## Conflict of interest statement

Nothing declared.

## Acknowledgements

Many thanks to Rebecca M Johnson, Dr Chen Heu for formatting and editing suggestions, to Drs Brittany Dodson and Donghun Kim for insightful discussion, and the anonymous reviewers for constructive suggestions. Funding support from NSF/BIO grant 1645331.

## References

1. Eisen RJ, Eisen L, Ogden NH, Beard CB: **Linkages of weather and climate with *Ixodes scapularis* and *Ixodes pacificus* (Acari: Ixodidae), enzootic transmission of *Borrelia burgdorferi*, and Lyme disease in North America.** *J Med Entomol* 2016, **53**:250-261.
2. Estrada-Pena A, Ayllon N, De La Fuente J: **Impact of climate trends on tick-borne pathogen transmission.** *Front Physiol* 2012, **3**.
3. Ogden NH, Lindsay LR: **Effects of climate and climate change on vectors and vector-borne diseases: ticks are different.** *Trends Parasitol* 2016, **32**:646-656.
4. Sonenshine DE: **Range expansion of tick disease vectors in North America: implications for spread of tick-borne disease.** *Int J Environ Res Public Health* 2018, **15**.
- The author present a review of current research on potential implications of climate change for disease transmission by four of the most important vector species of ticks in the USA.
5. Jore S, Vanwambeke SO, Viljugrein H, Isaksen K, Kristoffersen AB, Woldehiwet Z, Johansen B, Brun E, Brun-Hansen H, Westermann S et al.: **Climate and environmental change drives *Ixodes ricinus* geographical expansion at the northern range margin.** *Parasites Vectors* 2014, **7**:11.
6. Nava S, Venzal JM, González-Acuña D, Martins TF, Guglielmone AA: **Chapter 1 — tick classification, external tick anatomy with a glossary, and biological cycles.** *Ticks of the Southern Cone of America*. Academic Press; 2017:1-23.
7. Klompen H, Grimaldi D: **First mesozoic record of a parasitiform mite: a larval Argasid tick in Cretaceous Amber (Acari: Ixodida: Argasidae).** *Ann Entomol Soc Am* 2001, **94**:10-15.
8. Dunlop JA, Garwood RJ: **Tomographic reconstruction of the exceptionally preserved Trigonotarbid Arachnid *Eophrynus prestvicii*.** *Acta Palaeontol Pol* 2014, **59**:443-454.
9. Morshed M, Li L, Lee M-K, Fernando K, Lo T, Wong Q: **A Retrospective Cohort Study of Tick Paralysis in British Columbia.** *Vector Borne Zoonotic Dis* 2017, **17**:821-824.
10. Mead P, Hinckley A, Hook S, Beard CB: **TickNET — a collaborative public health approach to tickborne disease surveillance and research.** *Emerg Infect Dis* 2015, **21**:1574-1577.
11. Brites-Neto J, Duarte KMR, Martins TF: **Tick-borne infections in human and animal population worldwide.** *Vet World* 2015, **8**:301-315.
12. Guglielmone AA, Robbins RG, Apanaskevich DA, Petney TN, Estrada-Peña A, Horak IG: *The Hard Ticks of the World*. Netherlands: Springer; 2014.
13. Dantas-Torres F, Chomel BB, Otranto D: **Ticks and tick-borne diseases: a One Health perspective.** *Trends Parasitol* 2012, **28**:437-446.

Although this is not a paper from the last 2 years, it is an important paper that highlights the importance of cross-disciplinary communication for tick-borne zoonoses, putting them in the context of One Health.

14. Simon JA, Marrotte RR, Desrosiers N, Fiset J, Gaitan J, Gonzalez A, Koffi JK, Lapointe F-J, Leighton PA, Lindsay LR *et al.*: **Climate change and habitat fragmentation drive the occurrence of *Borrelia burgdorferi*, the agent of Lyme disease, at the northeastern limit of its distribution.** *Evol Appl* 2014, **7**:750-764.
15. de la Fuente J, Waterhouse RM, Sonenshine DE, Roe RM, Ribeiro JM, Sattelle DB, Hill CA: **Tick genome assembled: new opportunities for research on tick-host-pathogen interactions.** *Front Cell Infect Microbiol* 2016, **6**.
16. Donohoe H, Pennington-Gray L, Omodior O: **Lyme disease: current issues, implications, and recommendations for tourism management.** *Tour Manag* 2015, **46**:408-418.
17. Gulia-Nuss M, Nuss AB, Meyer JM, Sonenshine DE, Roe RM, Waterhouse RM, Sattelle DB, de la Fuente J, Ribeiro JM, Megy K *et al.*: **Genomic insights into the *Ixodes scapularis* tick vector of Lyme disease.** *Nat Commun* 2016, **7**:10507.
18. Hinckley AF, Connally NP, Meek JL, Johnson BJ, Kemperman MM, Feldman KA, White JL, Mead PS: **Lyme disease testing by large commercial laboratories in the United States.** *Clin Infect Dis* 2014, **59**:676-681.
19. Fang R, Blanton LS, Walker DH: **Rickettsiae as emerging infectious agents.** *Clin Lab Med* 2017, **37**:383-400.  
 Much attention is focused on the role of ticks in Lyme disease, but there are other tickborne zoonoses that are emerging, some which have fatal outcomes. While both pathogenic and nonpathogenic *Rickettsia* species have co-evolved with arthropod hosts, ticks transmit the majority of pathogenic species. This paper provides an updated review of rickettsioses and the increase in the number of incidences worldwide.
20. Chen LH, Wilson ME: **Tick-borne rickettsiosis in traveler returning from Honduras.** *Emerg Infect Dis* 2009, **15**:1321-1323.
21. González-Acuña D, Beldoménico PM, Venzal JM, Fabry M, Keirans JE, Guglielmone AA: **Reptile trade and the risk of exotic tick introductions into southern South American countries.** *Exp Appl Acarol* 2005, **35**:335-339.
22. Mansfield KL, Jizhou L, Phipps LP, Johnson N: **Emerging tick-borne viruses in the twenty-first century.** *Front Cell Infect Microbiol* 2017, **7**.
23. Mihalca AD: **Ticks imported to Europe with exotic reptiles.** *Vet Parasitol* 2015, **213**:67-71.
24. Molaei G, Andreadis TG, Anderson JF, Stafford Iii KC: **An exotic hitchhiker: a case report of importation into Connecticut from Africa of the human parasitizing tick, *Hyalomma truncatum* (Acari: Ixodidae).** *J Parasitol* 2018 <http://dx.doi.org/10.1645/18-13>.
25. Bardosh KL, Ryan S, Ebi K, Welburn S, Singer B: **Addressing vulnerability, building resilience: community-based adaptation to vector-borne diseases in the context of global change.** *Infect Dis Poverty* 2017, **6**.
26. Godfrey ER, Randolph SE: **Economic downturn results in tick-borne disease upsurge.** *Parasites Vectors* 2011, **4**:35.
27. Corn JL, Mertins JW, Hanson B, Snow S: **First reports of ectoparasites collected from wild-caught exotic reptiles in Florida.** *J Med Entomol* 2011, **48**:94-100.
28. Cohen EB, Auckland LD, Marra PP, Hamer SA: **Avian migrants facilitate invasions of neotropical ticks and tick-borne pathogens into the United States.** *Appl Environ Microbiol* 2015, **81**:8366-8378.  
 This paper highlights the importance of global spread of ticks and TBD through migrating birds. The authors present data on predominantly neotropical birds migrating between Texas, and Central or South America, the tick diversity, and the pathogenic bacteria that were present in both birds and ticks. They estimate that between 4 and 39 million ticks are introduced each year on migrating birds.
29. La Sorte FA, Fink D, Blancher PJ, Rodewald AD, Ruiz-Gutierrez V, Rosenberg KV, Hochachka WM, Verburg PH, Kelling S: **Global change and the distributional dynamics of migratory bird populations wintering in Central America.** *Glob Change Biol* 2017, **23**:5284-5296.
30. Mukherjee N, Beati L, Sellers M, Burton L, Adamson S, Robbins RG, Moore F, Karim S: **Importation of exotic ticks and tick-borne spotted fever group rickettsiae into the United States by migrating songbirds.** *Ticks Tick-Borne Dis* 2014, **5**:127-134.
31. Wu X, Röst G, Zou X: **Impact of spring bird migration on the range expansion of *Ixodes scapularis* tick population.** *Bull Math Biol N Y* 2016, **78**:138-168.
32. Eisen RJ, Kugeler KJ, Eisen L, Beard CB, Paddock CD: **Tick-borne zoonoses in the United States: persistent and emerging threats to human health.** *ILAR J* 2017 <http://dx.doi.org/10.1093/ilar/ilx005>.
33. Merten HA, Durden LA: **A state-by-state survey of ticks recorded from humans in the United States.** *J Vector Ecol J Soc Vector Ecol* 2000, **25**:102.
34. Takken W, Knols BGJ: *Olfaction in Vector-Host Interactions.* Wageningen Academic Publ; 2010.
35. Gaither M, Schumacher M, Nieto N, Corrigan J, Murray H, Maurer M: **Where are the ticks? Solving the mystery of a tickborne relapsing fever outbreak at a youth camp.** *J Environ Health Denver* 2016, **78**:8-11.
36. Lubelczyk C, Cahill BK, Hanson T, Turmel J, Lacombe E, Rand PW, Elias SP, Smith JRP: **Tick (Acari: Ixodidae) infestation at two rural, seasonal camps in Maine and Vermont.** *J Parasitol* 2010, **96**:442-443.
37. Rosenberg R: **Vital signs: trends in reported vectorborne disease cases — United States and Territories, 2004–2016.** *MMWR Morb Mortal Wkly Rep* 2018, **67**.
38. Steere AC, Broderick TF, Malawista SE: **Erythema chronicum migrans and Lyme arthritis: epidemiologic evidence for a tick vector.** *Am J Epidemiol* 1978, **108**:312-321.
39. Steere AC, Strle F, Wormser GP, Hu LT, Branda JA, Hovius JWR, Li X, Mead PS: **Lyme borreliosis.** *Nat Rev Dis Primer* 2016, **2**:16090.
40. Bouchard C, Leighton PA, Beauchamp G, Nguon S, Trudel L, Milord F, Lindsay LR, Bélanger D, Ogden NH: **Harvested white-tailed deer as sentinel hosts for early establishing *Ixodes scapularis* populations and risk from vector-borne zoonoses in southeastern Canada.** *J Med Entomol* 2013, **50**:384-393.
41. Lieske DJ, Lloyd VK: **Combining public participatory surveillance and occupancy modelling to predict the distributional response of *Ixodes scapularis* to climate change.** *Ticks Tick-Borne Dis* 2018, **9**:695-706.
42. Nelder MP, Russell C, Lindsay LR, Dhar B, Patel SN, Johnson S, Moore S, Kristjansson E, Li Y, Ralevski F: **Population-based passive tick surveillance and detection of expanding foci of blacklegged ticks *Ixodes scapularis* and the Lyme disease agent *Borrelia burgdorferi* in Ontario, Canada.** *PLOS ONE* 2014, **9**:e105358.
43. Ripoche M, Gasmi S, Adam-Poupart A, Koffi JK, Lindsay LR, Ludwig A, Milord F, Ogden NH, Thivierge K, Leighton PA: **Passive tick surveillance provides an accurate early signal of emerging Lyme disease risk and human cases in Southern Canada.** *J Med Entomol* 2018 <http://dx.doi.org/10.1093/jme/tjy030>.  
 The authors present a comparative analysis of passive surveillance, active surveillance, and reported human cases of Lyme disease in Canada. They found that passive tick surveillance was correlated with tick abundance (risk of tick encounter), but also a good indicator of Lyme disease risk.
44. Aenishaenslin C, Bouchard C, Koffi JK, Ogden NH: **Exposure and preventive behaviours toward ticks and Lyme disease in Canada: results from a first national survey.** *Ticks Tick-Borne Dis* 2017, **8**:112-118.
45. Kolopack PA, Parsons JA, Lavery JV: **What makes community engagement effective? Lessons from the eliminate dengue program in Queensland, Australia.** *PLoS Negl Trop Dis* 2015, **9**:e0003713.

The authors describe how Eliminate Dengue (now World Mosquito Program) was able to successfully implement its strategies by engaging community members.

46. Kowalec M, Szewczyk T, Welc-Fale?ciak R, Siński E, Karbowski G, Bajer A: **Ticks and the city — are there any differences between city parks and natural forests in terms of tick abundance and prevalence of spirochaetes?** *Parasit Vectors* 2017, **10**:573.
  47. Jones EH, Hinckley AF, Hook SA, Meek JI, Backenson B, ● Kugeler KJ, Feldman KA: **Pet ownership increases human risk of encountering ticks.** *Zoonoses Public Health* 2018, **65**:74-79.
- The authors present data supporting the correlation between pet ownership and likelihood of tick contact using data obtained through TickNET from 2727 households in the USA, the multi-state surveillance network established by the CDC. Given that over 70% of US households have at least one pet, veterinary professionals can play a significant role in TBD prevention.
48. Forrester JD, Vakkalanka JP, Holstege CP, Mead PS: **Lyme disease: what the wilderness provider needs to know.** *Wilderness Environ Med* 2015, **26**:555-564.
  49. Li S, Juhász-Horváth L, Trájer A, Pintér L, Rounsevell MDA, Harrison PA: **Lifestyle, habitat and farmers' risk of exposure to tick bites in an endemic area of tick-borne diseases in Hungary.** *Zoonoses Public Health* 2018, **65**:e248-e253.
  50. De Keukeleire M, Vanwambeke SO, Somassè E, Kabamba B, Luyasu V, Robert A: **Scouts, forests, and ticks: impact of landscapes on human-tick contacts.** *Ticks Tick-Borne Dis* 2015, **6**:636.
  51. Stott RE, Richards SL, Balanay JAG, Martin GL: **Prevention of tick exposure in environmental health specialists working in the Piedmont Region of North Carolina.** *J Environ Health Denver* 2016, **78**:E1-E7.
  52. Rabinowitz PM, Natterson-Horowitz BJ, Kahn LH, Kock R, Pappaioanou M: **Incorporating one health into medical education.** *BMC Med Educ* 2017, **17**.
  53. Elchos BN, Goddard J: **Implications of presumptive fatal Rocky Mountain spotted fever in two dogs and their owner.** *J Am Vet Med Assoc* 2003, **223**:1450-1452.
  54. Gibbs EPJ: **The evolution of One Health: a decade of progress and challenges for the future.** *Vet Rec* 2014, **174**:85-91.
  55. Bombara CB, Dürr S, Machovsky-Capuska GE, Jones PW, Ward MP: **A preliminary study to estimate contact rates between free-roaming domestic dogs using novel miniature cameras.** *PLOS ONE* 2017, **12**.
  56. Davies S, Abdullah S, Helps C, Tasker S, Newbury H, Wall R: **Prevalence of ticks and tick-borne pathogens: Babesia and Borrelia species in ticks infesting cats of Great Britain.** *Vet Parasitol* 2017, **244**:129-135.
  57. Rodríguez-Vivas RI, Apanaskevich DA, Ojeda-Chi MM, Trinidad-Martínez I, Reyes-Novelo E, Esteve-Gassent MD, Pérez de León AA: **Ticks collected from humans, domestic animals, and wildlife in Yucatan, Mexico.** *Vet Parasitol* 2016, **215**:106-113.
  58. Scott JD, Anderson JF, Durden LA, Smith ML, Manord JM, Clark KL: **Prevalence of the Lyme disease Spirochete, Borrelia burgdorferi, in Blacklegged Ticks, Ixodes scapularis at Hamilton-Wentworth, Ontario.** *Int J Med Sci* 2016, **13**:316-324.
  59. Shannon AB, Rucinsky R, Gaff HD, Brinkerhoff RJ: **Borrelia miyamotoi, other vector-borne agents in cat blood and ticks in Eastern Maryland.** *EcoHealth* 2017, **14**:816-820.
  60. Burroughs JE, Thomasson JA, Marsella R, Greiner EC, Allan SA: **Ticks associated with domestic dogs and cats in Florida, USA.** *Exp Appl Acarol* 2016, **69**:87-95.
  61. Day MJ: **One health: the importance of companion animal vector-borne diseases.** *Parasit Vectors* 2011, **4**:49.
  62. Cortinas R, Spomer SM: **Occurrence and county-level distribution of ticks (Acari: Ixodoidea) in Nebraska using Passive Surveillance.** *J Med Entomol* 2014, **51**:352-359.
  63. Koffi JK, Leighton PA, Pelcat Y, Trudel L, Lindsay LR, Milord F, Ogden NH: **Passive surveillance for I. scapularis ticks: enhanced analysis for early detection of emerging Lyme disease risk.** *J Med Entomol* 2012, **49**:400-409.
  64. Barrett AW, Noden BH, Gruntmeir JM, Holland T, Mitcham JR, Martin JE, Johnson EM, Little SE: **County scale distribution of Amblyomma americanum (Ixodida: Ixodidae) in Oklahoma: addressing local deficits in tick maps based on passive reporting.** *J Med Entomol* 2015, **52**:269-273.
  65. Abdullah S, Helps C, Tasker S, Newbury H, Wall R: **Ticks infesting domestic dogs in the UK: a large-scale surveillance programme.** *Parasites Vectors* 2016, **9**:391.
  66. Jameson LJ, Medlock JM: **Tick surveillance in Great Britain.** *Vector Borne Zoonotic Dis Larchmt N* 2011, **11**:403-412.
  67. Fernández-Ruvalcaba M, Preciado-De La Torre JF, Córdoba-Juarez G, García-Vázquez Z, Rosario-Cruz R, Saltijeral-Oaxaca J: **Animal bait effect on the recovery of Boophilus microplus larvae from experimentally infested grass in Morelos, Mexico.** *Parasitol Latinoam* 2003, **58**.
  68. Gherman CM, Mihalca AD, Dumitrache MO, Györke A, Oroian I, Sandor M, Cozma V: **CO<sub>2</sub> flagging — an improved method for the collection of questing ticks.** *Parasites Vectors* 2012, **5**:125.
  69. Rynkiewicz EC, Clay K: **Tick community composition in Midwestern US habitats in relation to sampling method and environmental conditions.** *Exp Appl Acarol* 2014, **64**:109-119.
  70. Tulloch JSP, McGinley L, Sanchez-Viscaino F, Medlock JM, Radford AD: **The passive surveillance of ticks using companion animal electronic health records.** *Epidemiol Infect Camb* 2017, **145**:2020-2029.
  71. Beaujean DJMA, Gassner F, Wong A, Steenbergen JE, Crutzen R, Ruwaard D: **Education on tick bite and Lyme borreliosis prevention, aimed at schoolchildren in the Netherlands: comparing the effects of an online educational video game versus a leaflet or no intervention.** *BMC Public Health* 2016, **16**.
  72. Beaujean DJMA, Crutzen R, Gassner F, Ameling C, Wong A, van Steenbergen JE, Ruwaard D: **Comparing the effect of a leaflet and a movie in preventing tick bites and Lyme disease in The Netherlands.** *BMC Public Health* 2016, **16**:495.
  73. Hofhuis A, van de Kasstele J, Sprong H, van den Wijngaard CC, Harms MG, Fonville M, van Leeuwen AD, Simões M, van Pelt W: **Predicting the risk of Lyme borreliosis after a tick bite, using a structural equation model.** *PLOS ONE* 2017, **12**:e0181807.
  74. Koffi JK, Savage J, Thivierge K, Lindsay LR, Bouchard C, Pelcat Y, Ogden NH: **Evaluating the submission of digital images as a method of surveillance for Ixodes scapularis ticks.** *Parasitol Camb* 2017, **144**:877-883.
  75. Hall JL, Alpers K, Bown KJ, Martin SJ, Birtles RJ: **Use of mass-participation outdoor events to assess human exposure to tickborne pathogens.** *Emerg Infect Dis* 2017, **23**:463-467.
- Although dragging or flagging might identify high tick load areas, this does not directly correlate with high risk of tick bite and/or TBD transmission if the area is not commonly trafficked. The authors used a very creative approach for tickborne disease surveillance and TBD education. The authors attended at 2-day marathon in Scotland through tick-infested terrain and collected ticks from roughly 500 participants to assess the frequency of tick contact by a large number of humans, as well as the frequency of pathogen-infected ticks.
76. Lewis J, Boudreau CR, Patterson JW, Bradet-Legris J, Lloyd VK: **Citizen science and community engagement in tick surveillance — a Canadian case study.** *Healthcare* 2018, **6**:22.
  77. Eisen L, Eisen RJ: **Critical evaluation of the linkage between tick-based risk measures and the occurrence of Lyme disease cases.** *J Med Entomol* 2016 <http://dx.doi.org/10.1093/jme/tjw092>.
  78. Seifert VA, Wilson S, Toivonen S, Clarke B, Prunuske A: **Community partnership designed to promote Lyme disease prevention and engagement in citizen science.** *J Microbiol Biol Educ* 2016, **17**:63-69.
- The authors utilized a multi-pronged approach to Lyme disease prevention, citizen science, and STEM education. The citizen scientists were high school students who, in addition to learning how to develop a hypothesis and design experiments, also learned basic tick bite prevention strategies.



79. Funk C, Gottfried J, Mitchell A: **Science news and information today**. *Pew Res Cent Journal Proj* 2017.
80. Smith A, Monica A: **Social media use in 2018**. *Pew Res Cent Internet Sci Tech* 2018.
81. Brownell SE, Price JV, Steinman L: **Science communication to the general public: why we need to teach undergraduate and graduate students this skill as part of their formal scientific training**. *J Undergrad Neurosci Educ* 2013, **12**:E6-E10.

This is another older, but relevant opinion piece which emphasized 5 years previously how important it is for scientists to be formally trained in effective scientific communication to lay audiences. The health field has begun to take these suggestions to heart, but the vector community

and entomologists as a whole could also benefit from formal science communication training.

82. National Science Board: **Chapter 7. science and technology: public attitudes and understanding**. *Science and Engineering Indicators 2018 (NSB-2018-1)*. National Science Foundation; 2018.

83. Office of HIV/AIDS and Infectious Disease Policy AS for H (ASH):  
• **21st Century Cures Act**. *HHS.gov* 2017.

Although Lyme disease was first described in the late 1970s, legislation to address the Lyme disease epidemic in the United States was only recently introduced in 2015, and again in 2016. In December of 2017, Department of Health and Human Services was authorized by The *21st Century Cures Act* to form a Tick-borne Disease Working Group.